# **Out of Hours**

# Inadequate glycaemic control in LMIC:

# health system failures in Peru



Photo: Socios en Salud, Sucursal Perú ©SES 2014.

Worldwide, 80% of patients with diabetes reside in low- and middle-income countries (LMICs). These patients are at higher risk of having inadequate glycaemic control and developing serious complications from their diabetes when compared with patients from high-income countries.1 There are multiple causes explaining poor outcomes, one being the failure of healthcare systems to adapt to the necessary challenges of managing patients with chronic diseases.

In Peru's highly fragmented health system, most patients use free or low-cost public health services. In comparison with private healthcare services, those using the public healthcare system report lower satisfaction rates and longer waiting times. Nevertheless, it seems that neither public nor private health systems provide adequate diabetes prevention and management services.2 To illustrate this situation, we created the story of a fictional Peruvian patient (Lucho), who has diabetes, basing him on typical experiences collected from patients with chronic diseases using the Peruvian public health system.

Lucho lives in a peri-urban area, and uses his local community health centre (CHC). Lucky to have been diagnosed at an early stage of his illness, he still goes through many difficulties receiving adequate diabetic care. After receiving the diabetic diagnosis a nurse at the CHC informed him that neither the glucose-lowering medications nor the equipment needed to screen for diabetic complications were in stock in that CHC. Lucho was therefore referred to the outpatient clinic of a public hospital 2 hours away from his house by public transport. Once in the hospital, Lucho requested an appointment, which was scheduled for 10 days later. On the appointment date, he

waited 2 hours before seeing a physician with the consultation lasting 15 minutes.<sup>3</sup> The physician spent much of this time filling in long clinical forms, and little time talking directly with Lucho. With such limited physician-patient interaction, Lucho wasn't able to ask any of the many questions he considered important. At the end of the appointment, the physician gave him a prescription for metformin and captopril, and a list of required laboratory tests, with a return visit scheduled for 3 months later. Once out of the doctor's office. Lucho went to the hospital's laboratory, where he was told to return in 2 days to conduct the laboratory

Once home, Lucho felt stressed and had many unanswered questions about his illness, so he sought help at the nearest CHC. However, he received contradictory recommendations from the staff, not trained in diabetes management. Eventually, he stopped taking his prescribed medications, and resumed inadequate lifestyle behaviours, such as eating a high-carbohydrate diet and taking little physical activity.

Lucho also became an easy target for charlatans promising him a quick cure for cash (a common scenario). Not surprisingly, his diabetes condition only worsened.

The Peruvian public health system is characterised by serious weaknesses in its infrastructure. For example, sometimes only a limited supply of drugs are available, there is a lack of basic laboratory tests such as glucose tests and a lack of trained health professionals. In addition, the fact there are few proper patient education programmes for chronic diseases plays its part.

One possible explanation for these deficiencies is that the healthcare system in Peru, as many healthcare systems in other LMIC, remains largely focused on the management of communicable diseases and prenatal care.4

In the context of ongoing demographic and epidemiologic transitions, many countries are facing growth in their elderly populations and an increase of chronic diseases burden. LMIC like Peru thus have to make radical changes to their public health systems. They must recognise chronic disease as a health priority and adopt a patient-centred approach ensuring that patients like Lucho receive the proper diabetic control, education, and social support required to meet their specific health

# ADDRESS FOR CORRESPONDENCE

#### J Jaime Miranda

CRONICAS — Centro de Excelencia en Enfermedades Crónicas, Universidad Peruana Cayetano Heredia, Av. Armendáriz 497, 2do Piso, Miraflores, Lima 18, Peru.

## E-mail: jaime.miranda@upch.pe

# Alvaro Taype-Rondan,

Research Fellow, CRONICAS Centre of Excellence in Chronic Diseases, Universidad Peruana Cayetano Heredia, Lima, Peru.

#### María Lazo-Porras.

Research Fellow, CRONICAS Centre of Excellence in Chronic Diseases, Universidad Peruana Cayetano Heredia, Lima, Peru.

#### Miguel Moscoso-Porras,

Research Fellow, CRONICAS Centre of Excellence in Chronic Diseases. Universidad Peruana Cavetano Heredia, Lima, Peru.

## Marcia Moreano-Sáenz,

Resident in Family Medicine, School of Medicine, Universidad Peruana Cayetano Heredia, Lima, Peru.

## J Jaime Miranda.

Director, CRONICAS Centre of Excellence in Chronic Diseases, Universidad Peruana Cayetano Heredia; and Research Professor, Department of Medicine, School of Medicine, Universidad Peruana Cayetano Heredia, Lima, Peru.

DOI: 10.3399/bjgp16X684541

# REFERENCES

- Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. Diabetes Res Clin Pract 2011; 94(3): 311-321.
- 2. Lopez Stewart G, Tambascia M, Rosas Guzmán J, et al. Control of type 2 diabetes mellitus among general practitioners in private practice in nine countries of Latin America. Rev Panam Salud Pública 2007; 22(1): 12-20.
- 3. Informatica INdEe. Encuesta nacional de Satisfacción de Usuarios del Aseguramiento Universal en Salud 2014. Lima: INEI, 2014. (National satisfaction survey in users of the universal health insurance 2014).
- Di Cesare M, Khang YH, Asaria P, et al. Inequalities in non-communicable diseases and effective responses. Lancet 2013; 381(9866): 585-597.
- 5. Atun R. Transitioning health systems for multimorbidity. Lancet 2015; 386(9995):